
Typically, <10 birds and more infrequently between 10 and 20 individuals of the species Redshank, Curlew, Turnstone, and Oystercatcher and more occasionally Dunlin recorded feeding on the mudflat in this area. The mudflat in this area regularly supports flocks <100 Black-tailed Godwit as well as Turnstone, Curlew, Dunlin (<50 birds) and lower numbers (<10 birds) of Oystercatcher, Redshank and Shelduck) Sea defences in Sector C are regularly used through the tide by individuals or small flocks of Turnstone (typically < 20 to 30 birds throughout sector). Shelduck, Redshank and Knot. .25 0.125 0.25



0

0.5



Immingham Green Energy Terminal

#### CLIENT

## **Associated British Ports**

## CONSULTANT

AECOM Limited 5th Floor 2 City Walk Leeds, LS11 9AR www.aecom.com

#### LEGEND



Site Boundary

Coastal Waterbirds

#### NOTES

Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS User Community.

## ISSUE PURPOSE

FINAL

## PROJECT NUMBER

# 60673509

FIGURE TITLE

The broad distribution of coastal waterbirds in Sector C

### FIGURE NUMBER

Figure 10.4